

B. Tech I Year Examinations, May/June -2012
MATHEMATICAL METHODS
 (Common to EEE, ECE, CSE, EIE, BME, IT, ETM, ECC, ICE)

Time: 3 hours

Max. Marks: 75

Answer any five questions
 All questions carry equal marks

1. a) Reduce the matrix into normal form, find its rank.

$$\begin{bmatrix} 2 & 3 & -1 & -1 \\ 1 & -1 & -2 & -4 \\ 3 & 1 & 3 & -2 \\ 6 & 3 & 0 & -7 \end{bmatrix}$$

- b) Find the values of 'a' and 'b' for which the equations, $x + y + z = 3$, $x + 2y + 2z = 6$, $x + 9y + az = b$ have
- No solution
 - A unique solution
 - Infinite number of solutions.

[8+7]

2. a) Find the Eigen values and the corresponding eigen vectors of the matrix

$$\begin{bmatrix} 6 & -2 & 2 \\ -2 & 3 & -1 \\ 2 & -1 & 3 \end{bmatrix}$$

- b) If λ is an Eigen value of a non-singular matrix A, then S.T. $\frac{|A|}{\lambda}$ is an Eigen value of Adj.A. [10+5]

3. Reduce the quadratic form $8x^2 + 7y^2 + 3z^2 - 12xy - 8yz + 4xz$ into a sum of squares by an orthogonal transformation and give the matrix of transformation. Also state the nature of the quadratic form. [15]

4. a) Find a real root of the equation $3x - \cos x - 1 = 0$ using Newton Raphson method.
 b) Find $f(1.6)$ using Lagranges formula from the following table. [8+7]

x	1.2	2.0	2.5	3.0
F(x)	1.36	0.58	0.34	0.20

5. a) Derive the normal equation to fit the parabola $y = a + bx + cx^2$.
 b) Given

x	1.0	1.1	1.2	1.3	1.4	1.5	1.6
Y=f(x)	7.989	8.403	8.781	9.129	9.451	9.750	10.031

Find y^1 and y^{11} at $x = 1.2$.

[7+8]

6. Find $y(0.1)$ and $y(0.2)$ using Runge Kutta fourth order formula given that $\frac{dy}{dx} = x + x^2y$ and $y(0) = 1$. [15]

7. a) Obtain the Fourier series expansion of $f(x)$ given that $f(x) = (\pi - x)^2$ in $0 < x < 2\pi$ and deduce the value of $\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots = \frac{\pi^2}{6}$.

- b) Obtain Fourier cosine series for $f(x) = x \sin x$ $0 < x < \pi$ and show that $\frac{1}{1.3} - \frac{1}{3.5} + \frac{1}{5.7} - \frac{1}{7.9} + \dots = \frac{\pi - 2}{4}$. [7+8]

8. a) Solve $(x^2 - yz)p + (y^2 - zx)q = z^2 - xy$.

- b) Form the partial differential equations by eliminating the arbitrary functions

i) $z = f(x^2 + y^2)$

ii) $z = yf(x) + xg(y)$. [7+8]