

3E1416

B.Tech. (Sem. III) (Main/Back) Examination, January - 2012
Automobile Engg.
3AE6 Advanced Engineering Mathematics
 (Common for ME/PI)

Time : 3 Hours]

[Total Marks : 80
 [Min. Passing Marks : 24

Instructions to Candidates :

Attempt any five questions selecting one question from each unit. All questions carry equal marks. Schematic diagrams must be shown wherever necessary. Any data you feel missing suitably be assumed and stated clearly. Units of quantities used/calculated must be stated clearly.

Use of following supporting material is permitted during examination.
 (Mentioned in form No. 205)

1. _____ Nil _____

2. _____ Nil _____

UNIT - I

- 1 (a) Find the Fourier series of

$$f(x) = x^2, \quad -\pi < x < 0$$

$$= -x^2, \quad 0 < x < \pi$$

In the interval given.

6

- (b) Obtain the Fourier cosine series for

$$f(x) = x^2, \quad 0 \leq x < 1$$

$$= 1, \quad 1 \leq x < 2$$

5

- (c) Find the Fourier transform of

$$f(t) = \begin{cases} -(1+t), & -1 \leq t < 0 \\ t-1, & 0 < t \leq 1 \\ 0, & |t| > 1 \end{cases}$$

5

OR

- 1 (a) State and prove the convolution theorem for Fourier transforms.

2, 6



(b) Use Fourier transforms to solve :

$$\frac{\partial u}{\partial t} = c^2 \frac{\partial^2 u}{\partial x^2}, \quad -\infty < x < \infty, \quad t > 0 \quad \text{and}$$

$$u(x, 0) = e^{-2|x|}, \quad -\infty < x < \infty$$

8

UNIT - II

2 (a) Solve the steady state temperature distribution equation :

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0, \quad 0 < x < a, \quad 0 < y < b$$

subject to $u(x, 0) = f(x)$, $u(x, b) = g(x)$, $0 < x < a$ and

$$u(0, y) = 0, \quad u(a, y) = 0, \quad 0 < y < b.$$

Use separation of variables method.

8

(b) (i) Find the Laplace transform of $\sin(3t+2)$

4

(ii) Find the Inverse Laplace transform of $\frac{1}{s(s^2+9)}$.

4

OR

2 (a) Use Laplace transforms solve $\frac{d^2 y}{dt^2} + 5 \frac{dy}{dt} + 4y = e^{3t}$,

$$y(0) = 0 \quad \text{and} \quad y'(0) = 2.$$

8

(b) (i) Find the Laplace transform of $\frac{1}{t}(1 - \cos bt)$.

4

(ii) Obtain the Inverse Laplace transform of $\frac{1}{(s^2+9)^2}$.

4

UNIT - III

3 (a) Determine the analytic function $f(z) = u + iv$ where

$$u + v = e^x (\cos y + \sin y).$$

8



(b) Show that $u = (y^3 - 3x^2y)$ is harmonic. Find the corresponding conjugate function $v(x, y)$ so that $f(z) = u + iv$ is analytic. 4

(c) Under the mapping $f(z) = z^2$, find the image of the region bounded by the lines $x=1$, $y=1$ and $x+y=1$. 4

OR

3 (a) State and prove Cauchy's integral theorem. 1, 4

(b) Find the poles of the function $f(z) = \frac{e^z}{(z - \sin z)}$

Also, find the principal part in the Laurent expansion of $f(z)$ about $z=0$. 2, 3

(c) Use contour integration to evaluate $\int_0^{2\pi} \frac{d\theta}{(3 + 2\sin \theta)}$

6

UNIT - IV

4 (a) A string is stretched between the fixed points $(0, 0)$ and $(l, 0)$ and released from rest from the initial deflection given by

$$f(x) = \begin{cases} \frac{2k}{l}x, & 0 < x < \frac{l}{2} \\ \frac{2k}{l}(l-x), & \frac{l}{2} < x < l \end{cases}$$

Find the deflection of the string at any time t . 8

(b) Obtain the steady state temperature distribution in a semi-circular plate whose bounding diameter is kept at 0°C while the circumference is kept at 50°C . 8

OR

4 (a) Solve in series : $(1-x^2) \frac{d^2y}{dx^2} - 2x \frac{dy}{dx} + 6y = 0$ 8

(b) Express $f(x) = 4x^3 + 3x^2 + 2x - 6$ in terms of Legendre polynomials. 4



(c) Evaluate : $\int_0^a x J_0(rx) dx$

4

UNIT - V

5 (a) Given

x :	2.5	3.0	3.5	4.0	4.5	5.0
y :	24.145	22.043	20.225	18.644	17.262	16.047

Obtain y for (i) $x=2.6$ (ii) $x=3.7$ and (iii) $x=4.8$.
State the formulae used.

3×3=9

(b) Given the following data :

x	300	304	305	307
y	2.4771	2.4829	2.4843	2.4871

Use Lagrange interpolation formula to obtain y for $x=306$.

7

OR

5 (a) Find $\frac{dy}{dx}$ and $\frac{d^2y}{dx^2}$ from the following data :

x	1.00	1.05	1.10	1.15	1.20	1.25	1.30
y	1.0000	1.0247	1.0488	1.0723	1.0954	1.1180	1.1401

for $x=1.00$.

3, 3

(b) Evaluate numerically $\int_0^{\pi/2} \sqrt{(\cos \theta)} d\theta$.

6

(c) Show that $\mu \delta = \frac{1}{2} (\Delta + \nabla)$, with usual meanings for symbols used.

4

