| | UTI | P7-RM-6 | |
|------|-------|--|-----|
| n | 220 | P7-RM-6 P7-RM-6 P7-RM-6 P7-RM-6 P7-RM-6 P7-RM-6 LJ-10 | 448 |
| /11 | . 002 | 26.11.13 | 440 |
| | | (3 Hours) [Total Marks: | 100 |
| | | | |
| N.B. | |) Question no. 1 is compulsory. | |
| | (2) | Answer any four questions out of remaining six questions. | |
| | | | |
| 1. | (a) | Find fourier exansion of $f(x) = 4-x^2$ in the ingterval $(0, 2)$ | 5 |
| | (b) | Find the probability that atmost 5 defective fuses will be found in a box of 200 | 5 |
| | | fuses if experience shows that 2% of such fuses are defective. | |
| | (c) | Given $6y = 5x + 90$, $15x = 18y + 130$, $6x^2 = 16$. Find | 5 |
| | | (i) $\frac{1}{x}$ and $\frac{1}{y}$ (ii) rand (iii) $6y^2$ | |
| | | Solve the two dimensional heat equation $\frac{d^2u}{dx^2} + \frac{d^2u}{dy^2} = 0$ which satisfies the conditions | |
| | (d) | Solve the two dimensional heat equation $dx^2 dy^2$ which satisfies the conditions | 5 |
| | | $u(0,y)=u(\ell,y)=u(x,0)=0$ and $u(x,a)=\sin\frac{n\pi x}{\ell}$. | |
| | | $u(0,y)=u(x,0)=0$ and $u(x,a)=\sin\frac{1}{\ell}$. | |
| 2. | (a) | Obtain fourier series for $f(x) = x - x^2$, $-\pi < x < \pi$ Hence deduce that | 7 |
| | | | |
| | | $\frac{\pi^2}{12} = \frac{1}{12} - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} + \dots$ | |
| | (b) | 1 2 3 4 | 7 |
| | (0) | show three or five? | |
| | (c) | A continuous random variable X has p.d.f. $f(x) = kx^2 e^{-x}$, $x \ge 0$. Find K, mean and | 6 |
| | | variance. | |
| | | | |
| 3. | (a) | Using normal distribution find the probability that in a group of 100 persons there | 7 |
| | | will be 55 males assuming that the probability of a person being male is $\frac{1}{2}$. | |
| | (b) | Derive wave equation for vibration of string | 7 |
| | (c) | Obtain fourier expansion of $f(x) = \sin x$ in the interval $(-\ell, \ell)$ where a is not an | 6 |
| | | integer. | |
| | | | |
| 4. | (a) | Calculate correlation coefficient from the foll: data | 7 |
| | | x : 23 27 28 29 30 31 33 35 36 39 | |
| | | y: 18 22 23 24 25 26 28 29 30 32 | |
| | (b) | A die was thrown 132 times and the foll: frequencies were observed: | 7 |
| | | No. obtained : 1 2 3 4 5 6 | |
| | | Frequency: 15 20 25 15 29 28 | |

Test the hypothesis that the die is unbiased.

[TURN OVER

(c) Obtain complex form of fourier series for $f(x) = \cosh 3x + \sinh 3x$ in (-3, 3).

6

7

6

7

6

- (a) A homogeneous rod of conducting material of length ℓ has ends kept at zero temperature
 and the temperature at the centre is T and falls uniformly to zero at the two ends.
 Find the temperature u(x, t) at any time.
 - (b) Obtain half range sine series for f(x) when

 $f(x) = x, \qquad 0 < x < \frac{\pi}{2}$

 $= \pi - x$, $\frac{\pi}{2} < x < \pi$ Hence deduce $\frac{\pi^2}{8} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots$

(c) Two independent samples of sizes 8 and 7 gave the foll: results

Sample 1: 19 17 15 21 16 18 16 14

Sample 2 : 15 14 15 19 15 18 16

is the difference between the sample means significant?

6. (a) Find the expansion of $f(x) = x(\pi - x)$, $0 < x < \pi$ as a half range cosine series. 7

Hence S. T

(i) $\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6}$ (ii) $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} = \frac{\pi^2}{12}$

- (b) The diameter of a semicircular plate of radius a is kept at 0° C and the temperature at the semicircular boundary is T° C. Find the steady state temperature $u(r, \theta)$.
- (c) The average of marks scored by 32 boys is 72 with standard deviation 8 while that of 36 girls is 70 with standard deviation 6. Test at 1% level of significance. Whether the boys perform better than the girls?
- 7. (a) S. T the functions $f_1(x) = 1$, $f_2(x) = x$ are orthogonal on (-1, 1). Determine the constants a and b such that the function $f_3(x) = -1 + ax + bx^2$ is orthogonal to both f_1 and f_2 on that internal.
 - (b) Find fourier integral representation of

f(x) = x, 0 < x < a= 0, x > a, f(-x) = f(x)

(c) If u = x-y, v = x+y and if x, y are uncorrelated,

P.T r uv = $\frac{6x^2 - 6y^2}{6x^2 + 6y^2}$